e-ISSN 2231 – 363X Print ISSN 2231 – 3621



Asian Journal of

PHARMACEUTICAL RESEARCH

Journal homepage: - www.ajprjournal.com

FORMULATION & EVALUATION OF MOUTH DISSOLVING TABLETS CONTAINING LOSARTAN POTASSIUM USING NATURAL SUPERDISINIGRANTS

Bhimi Kumari*, Abhishek Soni, Shivali Singla, Sachin Goyal, Samriti Thakur, Sunidhi Mahant

School of Pharmacy, Abhilashi Universitry, Mandi, Himachal Pradesh, India.

ABSTRACT

Losartan potassium used as antihypertensive drug but it goes under first pass metabolism due do which it has low bioavailability. Fast onset of action is major concern in the management of hypertension. Therefore, the aim of this study was to formulate mouth dissolving tablet of losartan potassium to improve its bioavailability, to achieve fast onset of action and increase patient compliance. Mouth dissolving tablets were prepared by direct compression method using natural super disintegrating agent (banana powder & apple pectin) and evaluated for precompression parameters and post compression parameters such as appearance, dimensions, hardness, weight variation, friability, wetting time ,dispersion time, water absorption ratio, disintegration & dissolution study. According to results of optimized batches it has been concluded that formulation batch F9 was an ideal batch which contain banana powder (2.5%) & cross povidone (2.5%) showed least disintegration time that is 26 seconds & maximum drug releases of (99.68%) within 12 minutes and was best among all the formulations.

Key words: Losartan Potassium, Mouth dissolving tablets, super disintegrating agent.

INTRODUCTION

In spite of various route of drug administration, the oral route of drug administration is the easiest, most suitable and mostly used route of drug administration for systemic effects. However, it is presumed that 90% of drugs which are used for systemic effects are administered through the oral route of drug administration [1]. The oral dosage forms require water for drug administration and when there is no water available then patient trouble. Many patient find difficulty to swallow tablet and hard gelatin capsule, consequently they do not take medication as prescribed. It is estimated that 50% of the population is affected by this problem which result high incident of incompliance and ineffective therapy. The difficulty is experienced in particular by pediatric and geriatric patients, but it also applied to people who are ill in bed and those active working patients who are busy or traveling, especially those who have no access to water. For these reasons, tablets which can rapidly dissolve or disintegrate in the oral cavity have attracted a great deal of attention.

Rapidly dissolving or disintegrating tablets are not only indicated for people who have swallowing difficulties, but also are ideal for active people [2-5].

United States Food and Drug Administration (FDA) defined MDT as "A solid dosage form containing medicinal substance or active ingredient which disintegrates rapidly usually within a matter of seconds when placed upon the tongue". The disintegration time for MDT generally ranges from several seconds to about 3 minutes [6-7].

Due to the presence of superdisintegrants, it gets dissolved quickly, resulting in rapid absorption of drug which results rapid onset of action. In mouth dissolving tablets the absorption is taking place directly from mouth that why do not suffers from the first pass metabolism, so bioavailability of drug increases [8].

Losartan Potassium is an angiotensin II receptor antagonist and used as antihypertensive drug. The molecular weight of Losartan Potassium is 461.0g/mol,

Corresponding Author :- Bhimi Kumari Email:- bhimikaundal31@gmail.com

half-life is 1.5 to 2hr, and its bioavailability is 25%-35%. It gets metabolized mainly in the liver. Mouth dissolving tablets are soluble in saliva are absorbed from the mouth, pharynx and esophagus as the saliva passes down into stomach, thus enhance the bioavailability by avoiding first pass metabolism. Mouth dissolving tablets also leads to an increased patient compliance, and fast onset of action. Keeping all these factors in mind, it was considered appropriate to formulate mouth dissolving of Losartan Potassium [9].

MATERIEL & METHOD

Losartan Potassium was received as a gift sample from Sun Pharma Ltd, Banana, apple and mixed fruit flavor was purchased from local market,MCC was taken from pharmaceutics laboratory which was purchased from Molychem Ltd, Aspartame, zinc stearate and cross povidone were obtained from pharmaceutics laboratory & were purchased from fromLobachem, HPMC was obtained from pharmaceutical chemistry laboratory which was purchased from Qualikems .SLS was taken from main store and it was purchased from central drug house P. Ltd.

Preparation of Fast Dissolving Tablet

Fast dissolving tablet tablets containing 25 mg of Losartan Potassium were prepared by direct compression method using formula give in Table 1. The drug and excipients were passed through 60 mesh sieve ensure better mixing. Avicel and mannitol was used as a directly compressible diluent. The directly compressible mixtures were compressed using multi punch tableting machine fitted with 8 mm flat punches. Before compression, the surface of die and punch were lubricated with zinc stearate [10].

EVALUATION

Pre Compression Parameters Angle of repose

Angle of repose is defined as the maximum angle possible between the surface of a pile of the Powder and horizontal plane. The improper flow of powder is due to frictional force between the particles and these frictional forces are quantified by angle of repose [11, 12].

tan (Angle of repose) = h / rAngle of repose = tan⁻¹ (h/r) Where, h = height of pile r = radius of the base of pile

Bulk Density

Bulk density is defined as the mass of a powder divided by the bulk volume. The bulk density of a powder depends primarily on particle size distribution, particle shape, and cohesiveness of particles.

Mathematically it is defined as:

Bulk Density (ρ_b) = w/V_b

Where, w = mass of powder

 $V_b = bulk volume$

Tapped Density

Tapped density is the ratio of mass of powder to the tapped volume. It was determined by mechanically tapping the measuring cylinder and the volume was noted.

Tapped density $(\boldsymbol{\rho_t}) = w / V_t$

Where, w = mass of powder $V_t = bulk$ volume

Carr's Compressibility Index

It is also one of the simple methods to evaluate flow property of a powder by comparing the bulk density and tapped density. It is calculated by:

Carr's Index = Tapped Density - Bulk Density Bulk Density

Hausners Ratio

It indicates the flow properties of the powder and is measured by the ratio of tapped density to the bulk density.

Hausner's ratio = Tapped density/Bulk density

Lower Hausner ratio (<1.25) indicates better flow properties than higher ones (>1.25).

POST COMPRESSION PARAMETER Appearance

Twenty tablets of each formulation were taken to check any physical or surface roughness in the tablet formulation [13, 14].

Dimension

Thickness and diameter were measured using a calibrated barmier caliper. Five tablets of each formulation were picked randomly and dimensions determined.

Uniformity of Weight

The test was performed according to specifications given in the Indian Pharmacopoeia, 2007 on 20 tablets which was selected randomly. The maximum acceptable limit is $\pm 7.5\%$ deviation of not more than two of the individual mass from average mass and none deviates by more than twice this percentage.

Measurement of Tablet Friability

Tablet friability was measured using the Roche Friabilator according to I.P.The friability was determined by following formula:-

 $F = W_A - W_B / W_A .100$

Where F-Friability W_A-Initial weight (g)

 $W_{\rm B}$ - Final weight (g).

B-1 mai weight (g).

Limit of friability for tablets under 1% is acceptable.

Measurement of Tablet hardness

The crushing strength of tablets was measured by a Monsanto Hardness Tester.

Wetting Time

A piece of tissue paper was folded twice and placed in small petri dish containing 6 ml of phosphate buffer (pH 6.8) the tablet was placed on it and the time required for complete wetting of tablet was recorded.

Water Absorption Ratio

A piece of tissue paper folded twice was placed in a small petri dish containing 6 ml of water. A tablet was put on the paper and was allowed for complete wetting. The wetted tablet was then weighed. Water absorption ratio, R, was determined using following equation:- $R = (W_A - W_B) / W_B * 100$

Where, W_{B} -Weight of tablet before water absorption W_{A} -Weight of tablet after water absorption

In vitro disintegration time

The process of breakdown of a tablet into smaller particles is called as disintegration. The in-vitro disintegration time of a tablet was determined using disintegration test apparatus as per I.P. specifications. Place one tablet in each of the 6 tubes of the basket. Add a disc to each tube and run the apparatususing distilled water maintained at $37^{0}(\pm 2^{0}\text{C})$ as the immersion liquid. The assembly should be raised and lowered between 30 cycles per minute in the pH 7.4 maintained at $37^{0}(\pm 2^{0}\text{C})$. The time in seconds taken for complete disintegration of the tablet with no palpable mass remaining in the apparatus was measured and recorded.

In vitro dissolution studies

Dissolution rate was studied by using IP type-I apparatus (50 rpm) using 900ml of water as dissolution medium. Temperature of the dissolution medium was maintained at 37.5° . The bath liquid is kept in constant and smooth motion during the testing. The sample was withdrawn at a regular time interval, filtered and diluted with the medium if necessary. The absorbance of filtered solution was measured by UV spectrophotometric method at 250 nm and concentration of the drug was determined from standard calibration curve.

Drug Content

At random 20 tablets were weighed and powdered. The powder equivalent to 50 mg was weighed accurately and dissolved in 100 ml of phosphate buffer of pH 6.8. The solution was shaken thoroughly. The undissolved matter was removed by filtration through Whatman No.1 filter paper. Then the serial dilutions were carried out. The absorbance of the diluted solutions was measured at 250 nm. The concentration of the drug was computed from the standard curve of the Losartan potassium in phosphate buffer of pH 6.8.

RESULT & DISCUSSION Pre formulations studies

The results for characterization of blended powder

are shown in Table 7.The bulk density of blend varied between 0.574-0.649 g/cm³. The tapped density was found in the range of 0.746-0.666 g/cm³. By using these two density data, Hausner's ratio and compressibility index was calculated. The powder blends of all formulation had Hausner's ratio of less than 1.25 indicating good flow characteristics. Blends having value of compressibility index less than 25% were considered as free flowing ones. The values for compressibility index were found between 12.346- 13.868. The flow ability of the powder was also evidenced by the angle of repose. The angle of repose below 35^{0} ranges indicates good flow properties of powder. The angle of repose was found to be in range $27-34^{0}$.

The powder flow properties were analyzed. It was observed that all formulations showed good flow properties with Carr's index ranging from 11.66 to 15.00 and Hausner's ratio below 1.25 which indicated good compressibility and flowability.

Post Compression studies

Appearance

All the tablets were white in colour, flat in shape with smooth surface without any defects.

Uniformity of thickness

The diameters of all the formulations was almost uniform (7.7-7.8mm). Thickness of all the formulations was found to be within the range of 2.23mm to 2.53mm shown in Table 3(a).

Weight Uniformity

All the tablets passed weight variation test as the % weight variation was within the pharmacopoeial limits of $\pm 7.5\%$. It is related to tooling of the compression machine, head pressure, machine speed and flow properties of the powder.

Hardness

The hardness of the Mouth dissolving tablet was found in the range of 3.5 to 4.3 kg/cm^2 is given in Table 3 (a).

Friability

Friability was observed less than 1% represented in Table 3(a), indicated that Mouth Dissolving Tablets had a good mechanical resistance. It is designed to evaluate the ability of the tablet to withstand abrasion in packaging, handling and shipping.

Wetting time

The wetting time was rapid in all the formulations. Wetting is closely related to inner structure of tablets, this may be due to ability of swelling and also capacity of absorption of water. Among all the formulations F9 showed less wetting time. The result was shown in Table 3(b).

Water absorption ratio

The capacity of disintegrant to swell in presence of little amount of water were found to be in the range of 54-66 % as shown in table 3(b). The water absorption ratio that is the up taking of water was very fast and the ratio was found higher.

In-vitro Disintegration time

Apple pectin & Banana Powder when comes in contact with water they quickly wicks water into the tablet through capillary action to create internal pressure that disintegrates tablet. The disintegration time is shown in table 3 (b).

Drug Content

The drug content was found to be within the range of 98.38 to 98.99 indicating uniform distribution of drug in the formulated tablets as per pharmacopoeia specification. The Drug content of all formulations was given in figure 2.

In vitro drug release studies

The comparative drug release was shown in

Figure2(a), 2(b) and 2 (c). Formulations F1 containing superdisintegrant Banana (2.5%), containing F2 superdisintegrant (3.5%),containing Banana F3 (5%), F4 containing superdisintegrant Banana superdisintegrant apple pectin (2.5)%), F5 containing superdisintegrant apple pectin (5%), F6 containing superdisintegrant apple pectin (5%), F7 containing superdisintegrant banana powder (1.5%) & cross povidone (1%), F8 containing superdisintegrant banana powder (2%) & cross povidone (1.5%), F9 containing superdisintegrant banana powder (2.5%) & cross povidone (2.5%), F10 containing superdisintegrant apple pectin (1.5%) & cross povidone (1%), F11 V containing superdisintegrant apple pectin (2%) & cross povidone (1.5%) and F12 containing superdisintegrant apple pectin (2.5%) & cross povidone (2.5%) showed a release of 92.58%, 95.76%, 96.01%, 93.45%, 94.53%, 96.56%, 97.46%, 99.68%, 93.86%, 94.06% and 95.34% respectively within twelve minutes. Among all the formulations resultant formulation F9 showed best release of 99.68% within twelve minutes.



Ingredients		Formulation Code										
(mg)	F1	F2	F3	F4	F5	F6	F7	F8	F9	F10	F11	F12
Losartan Potasium	25	25	25	25	25	25	25	25	25	25	25	25
Banana Powder	5	7	10	-	-	-	-	-	-	-	-	-
Apple Pectin	-	-	-	5	7	10	-	-	-	-	-	-
Banana Powder + Crosspovidone	-	-	-	-	-	-	3+2	4+3	5+5	-	-	-
Apple Pectin + Cross Povidone	-	-	-	-	-	-	-	-	-	3+2	4+3	5+5
MCC	78.5	77.5	76	78.5	77.5	76	78.5	77.5	76	78.5	77.5	76
D-Mannitol	78.5	77.5	76	78.5	77.5	76	78.5	77.5	76	78.5	77.5	76
HPMC	7	7	7	7	7	7	7	7	7	7	7	7
SLS	2	2	2	2	2	2	2	2	2	2	2	2
Zinc stearate	2	2	2	2	2	2	2	2	2	2	2	2
Aspartame	1	1	1	1	1	1	1	1	1	1	1	1
Mixed Fruit flavour	1	1	1	1	1	1	1	1	1	1	1	1
Total	200	200	200	200	200	200	200	200	200	200	200	200

 Table 1. Formulations of Losartan Potassium Fast Dissolving Tablets Prepared by Direct Compression Method (1 tablet)

Table 2. Evaluation of Pre Compression Parameters

Sr.	Ecomoletion Code	Angle of	Bulk	Тар	Carr's	Hausner's
No.	Formulation Code	Repose	Density	Density	Index	Ratio
1.	F1	28.36 ± 0.44	0.457 ± 0.002	0.530 ± 0.003	13.807 ± 0.104	1.16 ± 0.001
2.	F2	28.06±0.68	0.468 ± 0.002	0.542 ± 0.004	13.642 ± 0.596	1.158 ± 0.007
3.	F3	29.53±0.66	0.479 ± 0.002	0.557 ± 0.004	14.044 ± 0.637	$1.163\pm.008$
4.	F4	$33.54{\pm}0.60$	0.472 ± 0.002	0.552 ± 0.004	14.421 ± 0.641	1.168 ± 0.008
5.	F5	29.24± .43	0.466 ± 0.002	0.530 ± 0.004	12.058 ± 0.616	$1.137\pm.007$
6.	F6	$25.94 \pm .70$	0.474 ± 0.003	0.549 ± 0.004	13.763 ± 0.683	1.159 ± 0.009
7.	F7	28.06 ± 0.25	0.472 ± 0.003	0.544 ± 0.004	13.339 ± 0.638	1.153 ± 0.008
8.	F8	26.1 ± 0.46	0.477 ± 0.002	0.552 ± 0.004	13.515 ± 0.646	1.156 ± 0.008
9.	F9	$24.06 \pm .98$	0.489 ± 0.003	0.552 ± 0.004	11.464 ± 0.689	$1.129\pm.008$
10.	F10	25.63 ±0.46	0.495 ± 0.004	0.560 ± 0.004	11.598 ± 0.723	1.131 ± 0.009
11.	F11	$25.79 \pm .26$	0.489 ± 0.003	0.573 ± 0.004	14.758 ± 0.082	1.173 ± 0.001
12.	F12	$25.79 \pm .26$	0.483 ± 0.003	0.563 ± 0.008	14.203 ± 0.756	1.165 ± 0.010

Table 3 (a). Evaluation of Post Compression Parameters

Formulation	Thickness	Diameter	Hardness	Friability	
Code	(in mm)	(in mm)	(in kg/cm ²)	(in %age)	
F1	2.43 ± 0.057	7.76 ± 0.057	4.33 ± 0.152	0.585 ± 0.013	
F2	2.26 ± 0.152	7.76 ± 0.057	3.83 ± 0.057	0.728 ± 0.014	
F3	2.23 ± 0.115	7.83 ± 0.057	3.63 ± 0.152	0.590 ± 0.011	
F4	2.43 ± 0.115	7.83 ± 0.057	3.83 ± 0.305	0.585 ± 0.011	
F5	2.26 ± 0.057	7.86 ± 0.057	3.96 ± 0.208	0.448 ± 0.006	
F6	2.33 ± 0.057	7.83 ± 0.115	3.7 ± 0.2	0.583 ± 0.011	
F7	2.4 ± 0.173	7.83 ± 0.057	3.5 ± 0.4	0.555 ± 0.007	
F8	2.43 ± 0.115	7.83 ± 0.057	4.13 ± 0.208	0.588 ± 0.008	
F9	2.53 ± 0.115	7.83 ± 0.057	3.96 ± 0.208	0.440 ± 0.004	
F10	2.23 ± 0.115	7.83 ± 0.057	3.7 ± 0.1	0.492 ± 0.008	
F11	2.5 ± 0.1	7.76 ± 0.057	4.03 ± 0.23	0.540 ± 0.008	
F12	2.36 ± 0.057	7.83 ± 0.057	4.06 ± 0.152	0.689 ± 0.007	

Asian J. Pharm. Res. Vol 7, Issue 2, 62-67, 2017.

Formulation Code	Uniformity of Weight (in mg)	Wetting Time (in sec)	Water Absorption ratio (%)	Disintegration Time (in sec)
F1	198.85 ± 1.926	46	55.08	40.66 ± 1.254
F2	201.15 ± 1.814	41	58.56	39.66 ± 0.577
F3	200.6 ± 1.984	35	56.55	36.66 ± 1.527
F4	201.1 ± 1.832	53	59.56	51.33 ± 1.154
F5	200.6 ± 1.875	53	55.87	51.66 ± 1.527
F6	198.9 ± 1.889	52	60.46	54.66 ± 1.154
F7	198.75 ± 1.97	32	55.16	36.66 ± 1.154
F8	201.45 ± 1.904	35	61.42	28.66 ± 1.154
F9	200.4 ± 1.902	31	58.23	26.33 ± 1.152
F10	198.5 ± 1.986	54	55.23	41.33 ± 1.154
F11	198.85 ± 1.98	44	60.13	37.66 ± 0.57
F12	201.1 ±1.803	47	59.16	33.66 ± 1.54

Table 3 (b). Evaluation of Post Compression Parameters

CONCLUSION

The result was obtained and the conclusions drawn from the study are as follows: Losartan Potassium fast disintegrating tablet was prepared by direct compression method by using banana powder and apple pectin as superdisintegrants. Total 12 formulations were prepared and characterized. All the formulations showed fast disintegration of tablet. As the amount of banana powder and pectin increased in the formulation, there was decrease in disintegration of time. It was concluded that formulation F9 was the ideal batch from all the formulations which shows less disintegration time and maximum drug release within twelve minutes

ACKNOWLEDGEMENT Nil

IN11

CONFLICT OF INTEREST No interest

REFERENCES

- 1. Kishore HB, Sankar RK and Rao SB. Studies on Dissolution Rate of Paracetamol Tablets by Using Different Polymers. *Journal of Global Trends in Pharmaceutical Sciences*, 2, 2011, 1-10.
- 2. Yonezawa BY and Sunanda H. Rapidly disintegrating tablets prepared by the wet compression method: Mechanism and optimization. J. Pharm. Sci., 88(1), 1999, 1004-10.
- 3. Watanabe Y, Koizumi K, Zama Y, Matsumoto Y, Motsumoto M. New compressed tablet rapidly disintegrating in saliva in the mouth using crystalline cellulose and a disintegrant. *Bio. Pharm. Bull.*, 18(9), 1995, 1308-10.
- 4. Chang R, Guo X, Burnside B. A review of fast dissolving tablets. Pharm. Tech., 24(6), 2000, 52-4.
- 5. Abdelbary G, Prinderre P, Eouani C, Joachim J, Reynier JP. The preparation of orally disintegrating tablets using a hydrophilic waxy binder. *Int. J. Pharm.*, 278, 2004, 423-33.
- 6. Seong HJ, Yuuki T, Yourong F, Kinam P. Material properties for making fast dissolving tablets by a compression method. *J Mat Chem*, 18, 2008, 3527-3535.
- 7. Deshmukh KR, Patel V, Verma S, Pandey A, Dewangan P. A review on mouth dissolving tablet techniques. *Int J Research Ayu& Pharm*, 2(11), 2011, 66-74.
- 8. Dey P and Maiti S. Orodispersible tablets: A new trend in drug delivery. J. Nat. Sci. Biol. Med., 1, 2010, 2-5.
- 9. Jain CP, Naruka PS. Int J Pharmacy PharmSci, 1, 2009, 219-226.
- 10. Aitha S, Ayyappan T, Shanmugam S, Sundaramoorthy D and Vetrichelvan P. Optimization, Formulation and in-Vitro Evaluation of Mouth Dissolving Tablets of Levocetrizine Hydrochloride for the Treatment of Allergic Rhinitis. *J. Pharm. Sci. and Res*, 2, 2010, 555-561.
- 11. Subrahmanyam CVS. Textbook of physical pharmaceutics, Vallabhprakashan, 2nd edition 2000, 211-226.
- 12. Cooper J, Gun C. Powder Flow and Compaction. Inc Carter SJ, Eds. Tutorial Pharmacy. New Delhi: CBS Publication, 1986, 211-33.
- 13. Indian Pharmacopoeia, Government of India Ministry of Health & Family Welfare , Published by the Indian Pharmacopoeia Commission, Ghaziabad, 5th Edition, 2007.
- 14. Gohel MC, Bansal G and Bhatt N. Pharma Bio World, 3, 2005, 75-80.
- 15. Gohel M, Patel M, Amin A, Agrawal R, Dave R, Bariya N. AAPS. PharmSci-Tech, 5(3), 2005, 36.